



UNIVERSITI PUTRA MALAYSIA

**SEQUENCE AND FUNCTIONAL ANALYSES OF SALINITY
TOLERANCE GENES ISOLATED FROM THE MANGROVE PLANT,
ACANTHUS EBRACTEATUS (SEA HOLLY)**

NGUYEN PHUOC DANG.

FBSB 2005 3

**SEQUENCE AND FUNCTIONAL ANALYSES OF SALINITY TOLERANCE
GENES ISOLATED FROM THE MANGROVE PLANT,
Acanthus ebracteatus (SEA HOLLY).**

By

NGUYEN PHUOC DANG

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia,
in Fulfilment of the Requirements for the Degree of Doctor of Philosophy**

December 2005



Abstract of thesis presented to the Senate of Universiti Putra Malaysia in
fulfilment of the requirement for the degree of Doctor of Philosophy

**SEQUENCE AND FUNCTIONAL ANALYSES OF SALINITY TOLERANCE
GENES ISOLATED FROM THE MANGROVE PLANT,
Acanthus ebracteatus (SEA HOLLY).**

By

NGUYEN PHUOC DANG

December 2005

Chairman : Ho Chai Ling, PhD

Faculty : Biotechnology and Biomolecular Sciences

Salinity is a major abiotic stress that greatly affects plant growth and crop production. Most trees and crop plants are sensitive to salty conditions. Sodium ions are toxic to plants because of their adverse effects on potassium nutrition, cytosolic enzymes activities, photosynthesis and metabolism. Mangrove plants are good models to study plant tolerance to salinity as they possess salinity tolerance genes that allow them to survive under with high salinity conditions. The objectives of this study are to identify, isolate and characterize salinity tolerance genes from a mangrove plant, *Acanthus ebracteatus* using expressed sequence tag (EST) and bacterial functional assay approaches.

The leaves of *A. ebracteatus* were collected from the mangrove area at Morib, Selangor. Total RNA was isolated from the leaves of *A. ebracteatus*, and a cDNA library was constructed from cDNA fractionated between 500 to 5,000 bp. A total of eight hundred sixty four randomly selected clones were

isolated from the primary cDNA library from which 521 clones were sequenced. Among these ESTs, 138 of them were assembled into 43 contigs whereas 383 were singletons. A total of 349 of these ESTs showed significant homology to functional proteins and 18 % of them are particularly interesting as they correspond to genes involved in the stress response. Some of these clones, including mannitol dehydrogenase, plastidic aldolase, secretory peroxidase, ascorbate peroxidase, and vacuolar H⁺-ATPase, may be related to salinity tolerance mechanisms such as osmotic homeostasis, ionic homeostasis and detoxification.

In this study, a bacterial functional assay was also performed to identify cDNAs that confer salinity tolerance. A total of 120 salinity tolerant candidate genes from *A. ebracteatus* were isolated from 2 X YT medium supplemented with 400 mM NaCl and sequenced. Among these clones, 27 of them may be related to salinity tolerance such as manganese superoxide dismutase (Mn-SOD), putative salt tolerance protein, glutathione S-transferase, etc. The results showed that plants and bacteria may share some similar mechanisms for salinity tolerance.

A total of six cDNA clones from *A. ebracteatus* were fully sequenced and three of them were characterized by Southern hybridization and Northern hybridization. Clone A290 encoded a putative plastidic aldolase that may be involved in osmoprotection by converting triose phosphate into hexose. This gene was found to be expressed predominantly in the leaves of *A. ebracteatus*. There may be more than one family member of plastidic

aldolase in *A. ebracteatus*. Meanwhile, clone A303 was found to be a putative H⁺-ATPase, an enzyme known to play an important role in ion homeostasis, a salinity tolerance mechanism. This gene most probably exists as a single copy gene in *A. ebracteatus*. The expression of H⁺-ATPase was detected in all tissues of *A. ebracteatus*. Clone A325 encoded a putative monodehydroascorbate reductase which is involved in the detoxification mechanism. This gene was also expressed in all tissues and is most probably a single copy gene in the genome of *A. ebracteatus*.

Sequence analysis of the putative salinity tolerant cDNAs isolated by bacterial functional assay and ESTs suggested that the salinity tolerance mechanisms in *A. ebracteatus* may involve ion homeostasis, osmotic homeostasis, detoxification and other supporting mechanisms.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia
sebagai memenuhi keperluan untuk ijazah Doktor Falsafah

**ANALISIS JUJUKAN DAN FUNGSI GEN-GEN KETAHANAN TERHADAP
GARAM DARIPADA TUMBUHAN BAKAU,
Acanthus ebracteatus (SEA HOLLY).**

Oleh

NGUYEN PHUOC DANG

Desember 2005

Pengerusi : Ho Chai Ling, PhD

Fakulti : Bioteknologi dan Sains Biomolekul

Kegaraman (salinity) merupakan tekanan abiotik utama yang amat berkesan terhadap pertumbuhan pokok dan penghasilan tanaman. Kebanyakan pokok dan tanam-tanaman adalah sensitif terhadap keadaan yang masin. Ion natrium adalah toksik kepada tumbuhan disebabkan kesan yang buruk terhadap nutrisi potassium, aktiviti enzim sitosolik, fotosintesis dan metabolisme. Pokok bakau merupakan model yang baik untuk mengkaji ketahanan tumbuhan terhadap garam kerana mereka mempunyai gen ketahanan terhadap garam yang membolehkannya hidup di dalam kegaraman yang tingyi. Tujuan kajian ini adalah untuk mengenalpasti, memencilkan dan mencirikan gen ketahanan terhadap garam daripada pokok bakau, *Acanthus ebracteatus* dengan menggunakan 'expressed sequence tag' (EST) dan pendekatan esei bakteria berfungsi.

Daun *A. ebracteatus* telah dikumpulkan daripada kawasan hutan bakau di Morib, Selangor. RNA telah diekstrak daripada daun *A. ebracteatus* dan

perpustakaan cDNA telah dibina daripada fraksi cDNA di antara 500 ke 5000 bp. Sejumlah 864 klon telah dipilih secara rawak dan dipencilkan daripada perpustakaan cDNA primer di mana 521 klon telah dijuzuk. Di antara EST tersebut, 138 daripadanya wujud sebagai 43 kontig dan 383 yang selebihnya adalah singleton. Sebanyak 349 daripada EST ini menunjukkan homologi yang berkesan terhadap protein yang berfungsi dan 18 % daripadanya amat menarik kerana berhubungkait dengan gen-gen yang terlibat dengan tindak balas tekanan. Di antara klon-klon ini termasuk mannitol dehidrogenase, plastidik adolase, sekretori peroksidase, askorbat peroksidase, vakuolar H⁺-ATPase, yang mungkin berhubungkait dengan homeostasis osmotik, homeostasis ionik dan detoksifikasi.

Di dalam kajian ini, esei bakteria berfungsi dibuat untuk memencilkan cDNA yang mempunyai ketahanan terhadap garam. Sebanyak 120 gen yang menunjukkan ketahanan terhadap garam daripada *A. ebracteatus* telah dipencilkan daripada media 2 X YT yang mengandungi 400 mM NaCl dan analisis jujukan telah dibuat. Antara klon-klon ini, 127 daripadanya mungkin berhubungkait dengan protein ketahanan garam, glutathione S-transferase dan sebagainya. Keputusan menunjukkan bahawa tumbuhan dan bakteria berkongsi sesetengah mekanisme yang serupa di dalam ketahanan terhadap garam.

Sebanyak enam cDNA klon daripada *A. ebracteatus* telah jujuk sepenuhnya dan tiga klon cDNA telah dicirikan dengan menggunakan penghibridan 'Southern' dan 'Northern'. Klon A290 yang mengkodkan plastidik adolase,

mungkin terlibat di dalam 'osmoprotection' dengan mengubahkan trios fosfat kepada heksosa. Gen ini dizahirkan dengan banyaknya pada daun *A. ebracteatus*, dan didapati mungkin lebih daripada satu salinan dalam *A. ebracteatus* yang tergolong dalam keluarga yang sama. Manakala klon A303 merupakan putatif H^+ -ATPase, yang memainkan peranan penting di dalam homeostasis ion di dalam mekanisme toleransi terhadap garam. Gen ini berkemungkinan besar wujud sebagai satu salinan. Pengzahiran H^+ -ATPase telah dikesan pada semua tisu *A. ebracteatus*. Klon A325 yang mengekodkan putatif monodehidroaskorbat reduktase mungkin terlibat mekanisme nyah-toksik. Gen ini juga dizahirkan di dalam semua tisu dan berkemungkinan besar adalah salinan gen tunggal di dalam genom *A. ebracteatus*.

Analisa jujukan putatif cDNA yang berketahanan terhadap garam yang dipencilkan melalui esei bakteria berfungsi dan EST mencadangkan bahawa mekanisme ketahanan terhadap garam di dalam *A. ebracteatus* mungkin melibatkan hemoestasis ion, homeostasis osmotik, detoksifikasi dan mekanisme sokongan yang lain.

ACKNOWLEDGEMENTS

I would like to thank Dr. Ho Chai Ling for giving me the opportunity to work under her guidance for the last three years. Thanks for the encouragement. You have been an excellent mentor and your friendship as well as your guidance has been an honor to me.

I would like to express my sincere gratitude to my supervisors Assoc. Prof. Dr. Jennifer Ann Harikrishna, Assoc. Prof. Dr. Raha Abdul Rahim and Dr. Michael C.V.L.Wong for their guidance, support, advice and help throughout my thesis.

Many thanks to the laboratory members of Genetic laboratory, Faculty of Biotechnology and Biomolecular Sciences at Universiti Putra Malaysia: Mr. Ong, Choong, Lee, Swee Sen, Chan, Sudha, Seddon, Tony, Yeen Yee, Umami, Alfred, Joey, Sock Hwa, Tzer, Kam Lock, Siti and all the others, for the friendship, support, and encouragement.

I would like to thank Dr. Le Viet Dung, College of Agriculture and Cantho University, Viet Nam for providing the financial support during my study.

Many thanks to my friends Tung, Ky, Thuc, Mai Anh, Tam, Thach, Nghia in UPM for their support, and encouragement.

Last, but not least, many thanks go to my family for their love and support which have upheld me over the long years of my education.



I certify that an Examination Committee met on 14th December 2005 to conduct the final examination of Nguyen Phuoc Dang on his Doctor of Philosophy thesis entitled "Sequence and Functional Analyses of Salinity Tolerant Genes Isolated from the Mangrove Plant, *Acanthus ebractearus* (sea holly)" in accordance with Universiti Pertanian Malaysia (Higher Degree) Act 1980 and Universiti Pertanian Malaysia (Higher Degree) Regulations 1981. The Committee recommends that the candidate be awarded the relevant degree. Members of the Examination Committee are as follows:

Maziah Mahmood, PhD

Professor

Faculty of Biotechnology and Biomolecular Sciences
Universiti Putra Malaysia
(Chairman)

Siti Nor Akhmar Abdullah, PhD

Associate Professor

Faculty of Agriculture
Universiti Putra Malaysia
(Internal Examiner)

Norihan Mohd. Saleh, PhD


Associate Professor

Faculty of Biotechnology and Biomolecular Sciences
Universiti Putra Malaysia
(Internal Examiner)

Helen Nair, PhD

Professor

Faculty of Applied Sciences
Asian Institute of Medicine, Science and Technology
(External Examiner)


HASANAH MOHD. GHAZALI, PhD
Professor/Deputy Dean
School of Graduate Studies
Universiti Putra Malaysia

Date: 19 JAN 2006

This thesis submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfilment of the requirement for the degree of Doctor of Philosophy. The members of the Supervisory Committee are as follows:

Ho Chai Ling, PhD

Lecturer

Faculty of Biotechnology and Biomolecular Sciences

Universiti Putra Malaysia

(Chairman)

Jennifer Ann Harikrishna, PhD

Associate Professor

Malaysia University of Science and Technology

(Member)

Raha Abdul Rahim, PhD

Associate Professor

Faculty of Biotechnology and Biomolecular Sciences

Universiti Putra Malaysia

(Member)



AINI IDERIS, PhD

Professor/ Dean

School of Graduate Studies

Universiti Putra Malaysia

Date:

07 FEB 2006

DECLARATION

I hereby declare that the thesis is based on my original work except for quotations and citations which have been duly acknowledged. I also declare that it has not been previously or concurrently submitted for any other degree at UPM or other institutions.



NGUYEN PHUOC DANG

Date: 17 Jan 2006

TABLE OF CONTENTS

	Page
ABSTRACT	ii
ABSTRAK	v
ACKNOWLEDGEMENTS	viii
APPROVAL	ix
DECLARATION	xi
LIST OF TABLES	xvi
LIST OF FIGURES	xvii
LIST OF ABBREVIATIONS	xxi
 CHAPTER	
 1 INTRODUCTION	 1
 2 LITERATURE REVIEW	 4
2.1 Mangrove	4
2.1.1 Definition of Mangrove	4
2.1.2 Classification	5
2.1.3 The Ecology and Economical Role of Mangrove	7
2.1.4 Response of Mangrove to Salt	8
2.1.5 The Salt Tolerance Adaptation of Mangrove	10
2.2 <i>Acanthus ebracteatus</i> Vahl	12
2.3 Salt Stress in Plants	16
2.3.1 Effects of Salinity on Agriculture	16
2.3.2 Na ⁺ Toxicity in Plants	17
2.3.3 Sodium Influx in Plants	20
2.3.4 Sodium Efflux in Plants	22
2.4 Mechanism of Salt Tolerance	24
2.4.1 Homeostasis	25
2.4.2 Detoxification	30
2.4.3 Plant Growth Regulation	35
2.4.4 Salt Secretion by Salt Glands	36
2.5 Salt Tolerance Genes in Plants	38
2.6 RNA extraction	45
2.7 Expressed Sequence Tags (EST)	47
2.7.1 cDNA Library Generation and EST Sequencing	49
2.7.2 EST Clustering	50
2.7.3 Employing Bioinformatics Tools for Annotation of EST	51
2.8 Functional Assay	52
 3 MATERIALS AND METHODS	 56
3.1 Plant Materials-Sample Collection	56
3.2 Isolation of Total RNA	56
3.2.1 Method 1: Cesium chloride Method (Glišin <i>et al.</i> , 1974)	57

3.2.2	Method 2: Hexacetyltrimethyl ammonium bromide (CTAB method) (Apt <i>et al.</i> 1995)	58
3.2.3	Method 3: Modified Jill-Winter Method (Rochester <i>et al.</i> , 1986)	59
3.2.4	RNA Quantification	61
3.2.5	Formaldehyde Agarose Gel Electrophoresis	62
3.3	Isolation of mRNA from Total RNA	62
3.3.1	Isolation of mRNA	62
3.3.2	Quantification of mRNA	63
3.3.3	Ethidium bromide Plate Assay	64
3.4	Extraction of Genomic DNA	65
3.5	Construction of cDNA Library	66
3.5.1	First-Strand cDNA Synthesis	68
3.5.2	Second-Strand cDNA Synthesis	68
3.5.3	Blunting the cDNA Termini	69
3.5.4	Ligation with <i>EcoR</i> I Adapter	69
3.5.5	Phosphorylating the <i>EcoR</i> I End	70
3.5.6	Digestion with <i>Xho</i> I	70
3.5.7	Size Fractionation of cDNA	70
3.6	cDNA Purification	72
3.6.1	Ligating cDNA into the Uni-ZAP XR Vector	72
3.6.2	Packaging with Gigapack III Extract	73
3.7	Preparation of XL1 BlueMRF' Cells for Plating	72
3.8	Amplification of the cDNA Library	74
3.9	Random Selection of Plaques from Primary cDNA Library	75
3.9.1	<i>In vivo</i> Excision	75
3.9.2	Plasmid Preparation	76
3.10	PCR Amplification with Plasmid DNA	78
3.11	Bacterial Functional Assay	79
3.11.1	Screening for Salt Tolerant Candidates	79
3.11.2	Mass Excision	79
3.11.3	Plating	80
3.12	Sequence Analysis	81
3.13	Northern Blot Hybridization	83
3.14	Southern Blot Hybridization	84
3.15	Probe Preparation	87
3.16	Hybridization Analysis	87
4	RESULTS	90
4.1	RNA Extraction	90
4.2	Isolation of mRNA from Total RNA	94
4.3	Genomic DNA Extraction	94
4.4	Construction of cDNA Library	98
4.4.1	Quality of the Constructed cDNA Library	98
4.4.2	PCR-based Screening of <i>A. ebracteatus</i> cDNA clones	98
4.5	EST Analysis	101
4.6	Bacterial Functional Screening	112
4.7	Sequence Analyses and Molecular Characterization of	117

	Salt Tolerance Genes	
4.7.1	Sequence Analysis	117
4.7.2	Bacterial Functional Assay of Salt Tolerance Candidates	146
4.7.3	Molecular Analysis of Salt Tolerance Genes	146
5	DISCUSSION	156
5.1	Screening for Salt Tolerance Genes from <i>A. ebracteatus</i>	156
5.1.1	Expressed Sequence Tag Generation	156
5.1.2	Classification of ESTs	158
5.1.3	Putative Salt Tolerance Clones	161
5.2	Bacterial Functional Screening of Salt Tolerance Genes	168
5.3	Characterization of Putative Salt Tolerance Clones	171
5.3.1	Clone A290	173
5.3.2	Clone A303	175
5.3.3	Clone A325	177
6	CONCLUSION	182
	REFERENCES	186
	APPENDICES	214
	BIODATA OF THE AUTHOR	231

LIST OF TABLES

Table		Page
1	A comparison on the yield and purity of RNA from the three methods of RNA extraction from <i>A. ebracteatus</i> leaves	91
2	The yield and purity of RNA from modified Jill-Winter method from different tissues of <i>A. ebracteatus</i> .	93
3	ESTs from <i>A. ebracteatus</i> leaf cDNA library that may be related to salt tolerance	107
4	Putative function of some positive clones that may be related to salt tolerance.	114
5	Summary of 6 cDNA clones selected for full-length sequencing	118
6	The putative of candidate clones that may be related to salt stress tolerance.	163

LIST OF FIGURES

Figure		Page
1	Leaves of <i>Acanthus ebracteatus</i>	14
2	The flower of <i>A. ebracteatus</i>	15
3	Sodium ion uptake and extrusion at the plasma membrane. (reproduced from Blumwald <i>et al.</i> , 2000)	23
4	Pathway for reactive oxygen species (ROS) scavenging in plants. (reproduced from Asada, 2002)	34
5	The SOS pathway for salt tolerance. (reproduced from Zhu, 2002)	41
6	The cDNA synthesis flow chart. (source: cDNA synthesis kit instruction manual. Stratagene).	67
7	Northern blotting through gravitational transfer.	85
8	Formaldehyde agarose gel electrophoresis of RNA extracted from <i>A. ebracteatus</i> leaves using (A) CsCl method (lane 1), CTAB method (lane 2), and (B) Modified Jill-Winter method.	92
9	Formaldehyde agarose gel (1.5 % (w/v)) electrophoresis of RNA extracted from different tissues of <i>A. ebracteatus</i> using modified Jill-Winter method.	95
10	Estimation of mRNA concentration by using ethidium bromide plate assay.	96
11	Gel electrophoresis of DNA extracted from <i>A. ebracteatus</i> leaves.	97
12	Size fractionation of the double-stranded cDNA.	99
13	Amplification of cDNA inserts from randomly chosen clones of <i>A. ebracteatus</i> leaf cDNA library using T3 and T7 primers.	100
14	The size distribution of 731 cDNA inserts from the <i>A. ebracteatus</i> leaf cDNA library.	102
15	Distribution of ESTs from the <i>A. ebracteatus</i> leaf cDNA library according to BLASTX score.	104

16	The classification of ESTs from <i>A. ebracteatus</i> leaf cDNA library based on their putative functions.	106
17	The ESTs from <i>A. ebracteatus</i> leaf cDNA library match with genes of various plant species.	110
18	Classification of contigs according to the number of ESTs per contig.	111
19	The classification of cDNA sequences that conferred salt tolerance to <i>E. coli</i> based on their putative functions.	115
20	The cDNA sequences of <i>A. ebracteatus</i> leaf cDNA library from the bacterial functional screening with genes from various plant species.	116
21	The nucleotide and deduced amino acid sequence of clone A89. The asterisk indicates a stop codon.	121
22	Amino acid sequence alignment of A89 with secretory peroxidase from other species.	123
23	The nucleotide and deduced amino acid sequence of clone A163.	126
24	Amino acid sequence alignment of A163 with plastidic aldolase from other species.	128
25	The nucleotide and deduced amino acid sequence of clone A290.	131
26	Amino acid sequence alignment of A290 with plastidic aldolase from other species.	133
27	Amino acid sequence alignment of A163 with A290 amino acid sequences.	134
28	The nucleotide and deduced amino acids sequence of clone A303.	136
29	Amino acid sequence alignment of A303 with vacuolar H ⁺ -ATPase amino acid sequences from other species.	137
30	The nucleotide and deduced amino acids sequence of clone A310.	139
31	Amino acid sequence alignment of A310 with cytosolic ascorbate peroxidase from other species.	140

32	The nucleotide and deduced amino acids sequence of clone A325.	143
33	Amino acid sequence alignment of A325 with monodehydroascorbate reductase from other species.	145
34	Eight candidate clones from <i>A. ebracteatus</i> leaf cDNA library were examined on 2 X YT medium supplemented with or without 500 mM of NaCl.	147
35	Southern blot analysis with (A) <i>A. ebracteatus</i> genomic DNA (20 µg) digested with 6 restriction enzymes: 1- <i>Bam</i> H I, 2- <i>Eco</i> R I, 3- <i>Eco</i> R V, 4- <i>Hind</i> III, 5- <i>Not</i> I, 6- <i>Xba</i> I, M - λ <i>Pst</i> I marker. (B) Genomic southern blot analysis using A290 cDNA as probe.	150
36	Southern blot analysis with (A) <i>A. ebracteatus</i> genomic DNA (20 µg) digested with 6 restriction enzymes: 1- <i>Bam</i> H I, 2- <i>Eco</i> R I, 3- <i>Eco</i> R V, 4- <i>Hind</i> III, 5- <i>Not</i> I, 6- <i>Xba</i> I, M - λ <i>Pst</i> I marker. (B) Genomic Southern blot analysis using A303 as probe.	151
37	Southern blot analysis with (A) <i>A. ebracteatus</i> genomic DNA (20 µg) digested with 6 restriction enzymes: 1- <i>Bam</i> H I, 2- <i>Eco</i> R I, 3- <i>Eco</i> R V, 4- <i>Hind</i> III, 5- <i>Not</i> I, 6- <i>Xba</i> I, M - λ <i>Pst</i> I marker. (B) Genomic southern blot analysis using A325 as probe.	152
38	Northern blot analysis with (A) Expression of A290 (plastidic aldolase) in various tissues of <i>A. ebracteatus</i> . (B) As a control, the same blot was also hybridized with actin as loading control. (C) RNA (15 µg) was separated on a denaturing formaldehyde gel.	153
39	Northern blot analysis with (A) Expression of A303 (vacuolar-H ⁺ -ATPase) in various tissues of <i>A. ebracteatus</i> . (B) As a control, the same blot was also hybridized with actin as loading control. (C) RNA (15 µg) was separated on a denaturing formaldehyde gel.	154
40	Northern blot analysis with (A) Expression of A325 (monodehydroascorbate reductase) in various tissues of <i>A. ebracteatus</i> . (B) As a control, the same blot was also hybridized with actin as loading control. (C) RNA (15 µg) was separated on a denaturing formaldehyde gel.	155

LIST OF ABBREVIATIONS

α	alpha
β	beta
λ	lambda
$\times g$	gravitational acceleration
μg	microgram
μL	microliter
$^{\circ}C$	degree Centigrade
%	percentage
AMV	avian myeloblastosis virus
BLAST	Basic Local Alignment Search Tool
bp	base pairs
BSA	bovine serum albumin
Ca	calcium
cDNA	complementary DNA
CIP	calf intestinal phosphatase
Cl	chloride
cm	centimeter
CsCl	cesium chloride
CTAB	hexacetyltrimethyl ammonium bromide
dATP	2'-deoxy-adenosine-5'-triphosphate
dCTP	2'-deoxy-cytidine-5'-triphosphate
DEPC	diethyl pyrocarbonate
dGTP	2'-deoxy-guanosine-5'-triphosphate

DMSO	dimethylsulphonyl oxide
DNA	deoxyribonucleic acid
DNase	deoxyribonuclease
dNTPs	deoxynucleotides
ds	double-stranded
DTT	dithiothreitol
dTTP	thymidine-5'-triphosphate
EDTA	ethylenediaminetetraacetic acid
EtBr	ethidium bromide
g	gram
HCl	hydrochloric acid
HEPES	N-2-hydroxyethylpiperazine-N'-2 ethanesulfonic acid
IPTG	isopropyl- β -D-thiogalactoside
K	potassium
kb	kilo base-pair
L	liter
LB	Luria-bertani
LiCl	lithium chloride
M	molar
Mg	magnesium
MgCl ₂	magnesium chloride
MgSO ₄	magnesium sulfate
MOPS	3-(N-morpholino) propane-sulphonic acid
mL	milliliter
mM	millimolar

mRNA	messenger RNA
Na	sodium
NaCl	sodium chloride
NaOAc	sodium acetate
NaOH	sodium hydroxide
NCBI	National Center for Biotechnology Information
ng	nanogram
NH ₄ OAc	ammonium acetate
OD	optical density
ORF	open reading frame
PCR	polymerase chain reactions
PVP	polyvinylpyrrolidone
<i>pfu</i>	plaque forming units
ppm	part per million
RNA	ribonucleic acid
RNase A	ribonuclease A
ROS	reactive oxygen species
rpm	revolution per minute
RT	reverse transcriptase
SDS	sodium dodecyl sulphate
SOS	salt overly sensitive
TAE	tris acetate EDTA
TE	Tris-EDTA
U	unit
v/v	volume per volume

w/v	weight per volume
X-gal	5-bromo-4-chloro-3 indolyl- β -D-galactopyranoside

CHAPTER 1

INTRODUCTION

Salinity is one of the major abiotic stresses that affects plant growth and productivity globally. Salt stress can lead to changes in development, growth and productivity, and severe stress may threaten survival. High salinity causes both hyper osmotic and hyper ionic stress effects, and the consequence of these can be lethal to the plants. Therefore, a better understanding of the mechanisms that enable plants to adapt to salinity stress and to maintain growth will ultimately help in selection of stress tolerant cultivars for planting in saline soil.

In addition, due to the increased demand for food crops and plant products, the use of irrigated agriculture in the world has increased during the past 35 years (Chaturvedi, 2000). The rapid expansion in irrigation combined with the increase use waters containing high salt have led to the decrease in crop productivity, which is primarily due to salinity stress.

Mangroves represent the dominant soft bottom plant communities of the marine-terrestrial transition in tropical and subtropical regions. The mangrove species are members of terrestrial families that have adaptations to survive under conditions of high salinity, low oxygen and nutrient availability in the soil (Pernetta, 1993). Mangroves are divided into two distinct groups on the basis of their salt management strategies. One is “secreters” which have salt

glands or salt hairs and the other is “non secreters” lacking such morphological features for excretion of excess salt. The *Acanthus ebracteatus* is included in the first group.

Acanthus. spp is also known as Sea holly, holly mangrove, and “jeruju putih” (Malay). *A. ebracteatus* grows on the mud near the tide mark, often on mud lobsters mound. It can grow equally well under trees and in open areas, the plant can sometimes cover a large area and form thickets. The most striking feature of mangrove plant species is their ability to tolerate NaCl found in seawater up to the 500 mM level (Takemura *et al.*, 2000).

In order to elucidate salt tolerance mechanisms in higher plants, numerous key factors have been cloned such as late embryogenesis abundant protein (LEA) (Xu *et al.*, 1996), P5CS (Kishor *et al.*, 1995), DREB1A (Kasuga *et al.*, 1999), and AtNHX1 (Apse *et al.*, 1999). In contrast, the mechanisms that explain how plants can grow in saline conditions are still unclear. In order to grow under salinity stress, the mangrove plants must have acquired some proteins essential for salt tolerance mechanisms during their evolution. Recently, many reports addressing the mechanisms of mangrove plants at organ level were available (Werner and Stelzer, 1990). However, there were few reports about their mechanisms at molecular level because only a few model systems are available to analyze these mechanisms (Yamada *et al.*, 2002). Therefore, it is necessary to carry out molecular genetic studies on mangrove.